Author Index

Adeyemo, A.A., Oladosu, G.A. and Ayinla, A.O.
Growth and survival of fry of African catfish species, Clarias gariepinus Burchell,
Heterobranchus bidorsalis Geoffery and Heteroclarias reared on Moina dubia in
comparison with other first feed sources
Arai, K., see Kusunoki, T. et al.
Aulie, A., see Sohlberg, S. et al.
Ayinla, A.O., see Adeyemo, A.A. et al.
Bangen, M., Grave, K., Nordmo, R. and Søli, N.E.
Description and evaluation of a new surveillance programme for drug use in fish
farming in Norway
Banks, J.L.
Raceway density and water flow as factors affecting spring chinook salmon
(Oncorhynchus tshawytscha) during rearing and after release
Beacham, T.D., see Withler, R.E. et al.
Carrillo, M., see Piferrer, F. et al.
Castell, J.D., see Xu, X.L. et al.
Castell, L.L. and Mann, R.
Optimal staining of lipids in bivalve larvae with Nile Red
Chatain, B.
Abnormal swimbladder development and lordosis in sea bass (Dicentrarchus labrax)
and sea bream (Sparus auratus)
Chen, HY. and Tsai, JC.
Optimal dietary protein level for the growth of juvenile grouper, Epinephelus
malabaricus, fed semipurified diets
Clarke, W.C., see Whyte, J.N.C. et al.
Clement, S. and Lovell, R.T.
Comparison of processing yield and nutrient composition of cultured Nile tilapia
(Oreochromis niloticus) and channel catfish (Ictalurus punctatus)
Deguchi, Y., see Sugita, H. et al.
Divanach, P., see Kentouri, M. et al.
Donaldson, E.M., see Piferrer, F. et al.
Douillet, P.A. and Langdon, C.J.
Use of a probiotic for the culture of larvae of the Pacific oyster (Crassostrea gigas
Thunberg)
Drori, S., Ofir, M., Levavi-Sivan, B. and Yaron, Z.
Spawning induction in common carp (Cyprinus carpio) using pituitary extract or
GnRH superactive analogue combined with metoclopramide: analysis of hormone
profile, progress of oocyte maturation and dependence on temperature
Eid, A.E. and Ghonim, S.I.
Dietary zinc requirement of fingerling Oreochromis niloticus
Fracalossi, D.M. and Lovell, R.T.
Dietary lipid sources influence responses of channel catfish (Ictalurus punctatus) to
challenge with the pathogen Edwardsiella ictaluri 287

Ghonim, S.I., see Eid, A.E. et al.
Ginther, N.G., see Whyte, J.N.C. et al.
Goard, L.J., see Nell, J.A. et al.
González, S., see Vázquez, R. et al.
Grave, K., see Bangen, M. et al.
Gravel, M., see Sarac, H.Z. et al.
Guiral, D., see Konan-Brou, A.A. et al.
Harboe, T., Huse, I. and Øie, G.
Effects of egg disinfection on yolk sac and first feeding stages of halibut (Hippoglossus
hippoglossus L.) larvae
Harrison, P.J., see Thompson, P.A. et al.
Heasman, M.P., see Nell, J.A. et al.
Himmelman, J.H., see Lodeiros, C.J. et al.
Hirato, C., see Sugita, H. et al.
Huse, I., see Harboe, T. et al.
Jensen, J.O.T., see Whyte, J.N.C. et al.
Ji, W.J., see Xu, X.L. et al.
Kentouri, M., León, L., Tort, L. and Divanach, P.
Experimental methodology in aquaculture: modification of the feeding rate of the
gilthead sea bream Sparus aurata at a self-feeder after weighing
Konan-Brou, A.A. and Guiral, D.
Available algal biomass in tropical brackish water artificial habitats
Kuruma, A., see Sugita, H. et al.
Kusunoki, T., Arai, K. and Suzuki, R.
Production of viable gynogens without chromosome duplication in the spinous loach
Cobitis biwae
Langdon, C.J., see Douillet, P.A. et al.
Lee, CS., see Tamaru, C.S. et al.
León, L., see Kentouri, M. et al.
Levavi-Sivan, B., see Drori, S. et al.
Lodeiros, C.J. and Himmelman, J.H.
Relations among environmental conditions and growth in the tropical scallop Euvola
(Pecten) ziczac (L.) in suspended culture in the Golfo de Cariaco, Venezuela 345
Lovell, R.T., see Clement, S. et al.
Lovell, R.T., see Fracalossi, D.M. et al.
Mann, R., see Castell, L.L. et al.
McMeniman, N.P., see Sarac, H.Z. et al.
Montagnes, D.J.S., see Thompson, P.A. et al.
Mourente, G., see Vázquez, R. et al.
Murashige, R., see Tamaru, C.S. et al.
Nell, J.A., O'Connor, W.A., Heasman, M.P. and Goard, L.J.
Hatchery production for the venerid clam Katelysia rhytiphora (Lamy) and the Sydney
cockle Anadara trapezia (Deshayes)
Nordmo, R., see Bangen, M. et al.
Ohandarff T and Danshan I D
Oberdorff, T. and Porcher, J.P.
An index of biotic integrity to assess biological impacts of salmonid farm effluents on
An index of biotic integrity to assess biological impacts of salmonid farm effluents on receiving waters
An index of biotic integrity to assess biological impacts of salmonid farm effluents on receiving waters
An index of biotic integrity to assess biological impacts of salmonid farm effluents on receiving waters

Øie, G., see Harboe, T. et al.
Okada, R., see Sugita, H. et al.
Oladosu, G.A., see Adeyemo, A.A. et al.
Pedersen, H.P., see Staurnes, M. et al.
Piferrer, F., Carrillo, M., Zanuy, S., Solar, I.I. and Donaldson, E.M.
Induction of sterility in coho salmon (Oncorhynchus kisutch) by androgen immersion
before first feeding
Porcher, J.P., see Oberdorff, T. et al.
Rodríguez, A., see Vázquez, R. et al.
Rustad, T., see Staurnes, M. et al.
Sarac, H.Z., McMeniman, N.P., Thaggard, H., Gravel, M., Tabrett, S. and Saunders, J.
Relationships between the weight and chemical composition of exuvia and whole body
of the black tiger prawn, Penaeus monodon
Saunders, J., see Sarac, H.Z. et al.
Scarpa, J., Toro, J.E. and Wada, K.T. Direct comparison of six methods to induce triploidy in bivalves
Shaw, B.A., see Thompson, P.A. et al. Shearer, K.D.
Factors affecting the proximate composition of cultured fishes with emphasis on
salmonids
Sigholt, T., see Staurnes, M. et al.
Sin, F.Y.T., see Symonds, J.E. et al.
Sohlberg, S., Aulie, A. and Søli, N.E.
Temperature-dependent absorption and elimination of flumequine in rainbow trout
(Oncorhynchus mykiss Walbaum) in fresh water
Solar, I.I., see Piferrer, F. et al.
Søli, N.E., see Bangen, M. et al.
Søli, N.E., see Sohlberg, S. et al.
Staurnes, M., Sigholt, T., Pedersen, H.P. and Rustad, T.
Physiological effects of simulated high-density transport of Atlantic cod (Gadus
morhua)
Sugita, H., Kuruma, A., Hirato, C., Ohkoshi, T., Okada, R. and Deguchi, Y.
The vitamin-B ₁₂ -producing bacteria in the water and sediment of a carp culture pond 425
Suzuki, R., see Kusunoki, T. et al.
Symonds, J.E., Walker, S.P. and Sin, F.Y.T.
Electroporation of salmon sperm with plasmid DNA: evidence of enhanced sperm/
DNA association
Tabrett, S., see Sarac, H.Z. et al.
Tamaru, C.S., Murashige, R. and Lee, CS.
The paradox of using background phytoplankton during the larval culture of striped
mullet, Mugil cephalus L
Thaggard, H., see Sarac, H.Z. et al.
Thomas, D.H.L.
Socio-economic and cultural factors in aquaculture development: a case study from
Nigeria
Thompson, P.A., Montagnes, D.J.S., Shaw, B.A. and Harrison, P.J.
The influence of three algal filtrates on the grazing rate of larval oysters (Crassostrea
gigas), determined by fluorescent microspheres
Toro, J.E., see Scarpa, J. et al.
Tort, L., see Kentouri, M. et al.
Townsend, L.D., see Whyte, J.N.C. et al.
Tsai, JC., see Chen, HY. et al.

Vázquez, R., González, S., Rodríguez, A. and Mourente, G.
Biochemical composition and fatty acid content of fertilized eggs, yolk sac stage larvae
and first-feeding larvae of the Senegal sole (Solea senegalensis Kaup)273
Wada, K.T., see Scarpa, J. et al.
Walker, S.P., see Symonds, J.E. et al.
Whyte, J.N.C., Clarke, W.C., Ginther, N.G., Jensen, J.O.T. and Townsend, L.D.
Influence of composition of Brachionus plicatilis and Artemia on growth of larval
sablefish (Anoplopoma fimbria Pallas)
Withler, R.E. and Beacham, T.D.
Genetic variation in body weight and flesh colour of the coho salmon (Oncorhynchus
kisutch) in British Columbia
Xu, X.L., Ji, W.J., Castell, J.D. and O'Dor, R.K.
Influence of dietary lipid sources on fecundity, egg hatchability and fatty acid
composition of Chinese prawn (Penaeus chinensis) broodstock
Yaron, Z., see Drori, S. et al.
Zanuv S. see Piferrer F. et al.